

BEST AVAILABLE COPY

(86) Date de dépôt PCT/PCT Filing Date: 2000/03/28
(87) Date publication PCT/PCT Publication Date: 2000/10/19
(85) Entrée phase nationale/National Entry: 2001/10/01
(86) N° demande PCT/PCT Application No.: EP 2000/002711
(87) N° publication PCT/PCT Publication No.: 2000/061764
(30) Priorité/Priority: 1999/04/09 (199 16 140.2) DE

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/53, C12N 9/02, C12P 23/00
(71) Demandeur/Applicant:
BASF AKTIENGESELLSCHAFT, DE
(72) Inventeurs/Inventors:
SANDMANN, GERHARD, DE;
LINDEN, HARTMUT, DE
(74) Agent: ROBIC

(54) Titre : CAROTENE-HYDROXYLASE ET TECHNIQUE PERMETTANT DE PRODUIRE DES DERIVES DE XANTHOPHYLLE
(54) Title: CAROTENE HYDROXYLASE AND METHOD FOR PRODUCING XANTHOPHYLL DERIVATIVES

(57) Abrégé/Abstract:

The invention relates to proteins with an enzymatic activity for converting β -carotene into zeaxanthin or canthaxanthin into astaxanthin, to nucleic acids that code for these proteins, to nucleic acid constructs containing these nucleic acids, to genetically modified organisms in which the genetic modification causes or increases the gene expression of this nucleic acid compared with a wild type, and to methods for producing xanthophyll derivatives.

(51) Internationale Patentklassifikation ⁷: C12N 15/53, 9/02, C12P 23/00	A1	(11) Internationale Veröffentlichungsnummer: WO 00/61764 (43) Internationales Veröffentlichungsdatum: 19. Oktober 2000 (19.10.00)
(21) Internationales Aktenzeichen: PCT/EP00/02711 (22) Internationales Anmeldedatum: 28. März 2000 (28.03.00) (30) Prioritätsdaten: 199 16 140.2 9. April 1999 (09.04.99) DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): LINDEN, Hartmut [DE/DE]; St. Johannngasse 4, D-78462 Konstanz (DE). SANDMANN, Gerhard [DE/DE]; H. Steinhäuser Strasse 18, D-63065 Offenbach (DE). (74) Gemeinsamer Vertreter: BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE).	(81) Bestimmungsstaaten: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO Patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Veröffentlicht Mit internationalem Recherchenbericht.	

(54) Title: CAROTENE HYDROXYLASE AND METHOD FOR PRODUCING XANTHOPHYLL DERIVATIVES

(54) Bezeichnung: CAROTINHYDROXYLASE UND VERFAHREN ZUR HERSTELLUNG VON XANTHOPHYLLDERIVATEN

(57) Abstract

The invention relates to proteins with an enzymatic activity for converting β -carotene into zeaxanthin or canthaxanthin into astaxanthin, to nucleic acids that code for these proteins, to nucleic acid constructs containing these nucleic acids, to genetically modified organisms in which the genetic modification causes or increases the gene expression of this nucleic acid compared with a wild type, and to methods for producing xanthophyll derivatives.

(57) Zusammenfassung

Die vorliegende Erfindung betrifft Proteine, die eine enzymatische Aktivität zur Umwandlung von β -Carotin in Zeaxanthin oder Canthaxanthin in Astaxanthin aufweisen, Nukleinsäuren, die diese Proteine codieren, Nukleinsäurekonstrukte, enthaltend diese Nukleinsäuren, genetisch veränderte Organismen, wobei die genetische Veränderung die Genexpression dieser Nukleinsäure verglichen mit einem Wildtyp verursacht oder erhöht, sowie Verfahren zur Herstellung von Xanthophyllderivaten.

CAROTENE HYDROXYLASE AND METHOD FOR
PRODUCING XANTHOPHYLL DERIVATIVES

The present invention relates to proteins which have an enzymatic activity for converting β -carotene into zeaxanthin or canthaxanthin into astaxanthin, to nucleic acids which encode these proteins, to nucleic acid constructs comprising these nucleic acids, to genetically manipulated organisms where the genetic manipulation causes or increases the gene expression of this nucleic acid by comparison with a wild-type, and to processes for preparing xanthophyll derivatives.

- 10 Xanthophylls are oxygen-containing carotenoids of animal, plant or microbial origin. Xanthophylls such as lutein, zeaxanthin or astaxanthin are important additives in the human and livestock diet as pigmenting substances and precursors of vitamin A derivatives. In addition, xanthophylls have a health-promoting action such as enhancing the immune response and, by reason of their antioxidant properties, a cancer-preventing action, which makes their use as nutraceuticals of interest. An economic process for preparing xanthophylls and foodstuffs with an increased xanthophyll content are therefore of great importance. Particularly economic processes for preparing xanthophylls are biotechnological processes which make use of proteins and biosynthesis genes of xanthophyll biosynthesis from xanthophyll-producing organisms.

- 20 Prokaryotic β -carotene hydroxylases which catalyze the enzymatic conversion of β -carotene into zeaxanthin via β -cryptoxanthin, and the genes which encode these proteins are known from the bacteria *Erwinia uredovora* (Misawa et al., J. of Bacteriology 1990, 6704-6712; EP 393690 B1), *Erwinia herbicola* (WO 9113078), *Agrobacterium aurantiacum* (Misawa et al., J. of Bacteriology 1995, 6575-6584; EP 735 137 A1), *Alcaligenes sp. PC-1* (EP 735 137 A1), *Flavobacterium sp. strain R1534* (Pasamontes et al., Gene 1997, 185, 35-41; EP 747483 A2) and from the Cyanobacterium *Synechocystis sp. PCC6803* (Masamoto et al., Plant Cell Physiol. 1998, 39(5), 560-564).

- It is also known that the prokaryotic β -carotene hydroxylases from
30 *Agrobacterium aurantiacum*, *Alcaligenes* and *Erwinia uredovora* are additionally able to covert canthaxanthin via adonirubin in astaxanthin (Misawa et al., J. of Bacteriology 1995, 6575-6584; Fraser et al., J. Biol. Chem. 1997, 272, 6128-6135).

From eukaryotic sources, three plant β -carotene hydroxylases are known to catalyze the enzymatic conversion of β -carotene into zeaxanthin via β -cryptoxanthin. The corresponding cDNAs have been isolated from *Arabidopsis thaliana* (Cunningham et al, J. Biol. Chem. 1996, 271, 24349-24352, WO 9736998), and from *Capsicum annuum* L. (Bouvier et al., Biochimica et Biophysica Acta 1998, 1391, 320-328).

Genes of eukaryotic origin have the advantage over prokaryotic genes that they are expressed better in higher transgenic organisms such as plants. Nevertheless, there is still a need to improve and increase the xanthophyll productivity for an economic process for preparing xanthophyll derivatives or foodstuffs with an increased xanthophyll content by incorporating eukaryotic nucleic acids into organisms.

In addition, the appropriate eukaryotic β -carotene hydroxylases in the prior art have the disadvantage that they have only a narrow substrate range so that there is a build up of metabolic products which cannot be converted by the hydroxylases and may exert an inhibiting effect on the hydroxylases.

It is an object of the present invention to remedy the described deficiencies of the prior art and to provide a eukaryotic β -carotene hydroxylase with improved properties.

We have found that this object is achieved by a protein which has an enzymatic activity for converting β -carotene into zeaxanthin or canthaxanthin into astaxanthin, comprising the amino acid sequence SEQ ID NO. 2 or a sequence derived from this sequence by substitution, insertion or deletion of amino acids and having a homology of at least 50% at the amino acid level with the sequence SEQ ID NO.2.

Carotene hydroxylases hereinafter mean the proteins according to the invention, i.e. proteins which have an enzymatic activity for converting β -carotene into zeaxanthin or canthaxanthin into astaxanthin, comprising the amino acid sequence SEQ ID NO. 2 or a sequence derived from this sequence by substitution, insertion or deletion of amino acids and having a homology of at least 50% at the amino acid level with the sequence SEQ ID NO.2.

The amino acid sequence depicted in SEQ ID NO. 2 is derived from translation of the cDNA sequence depicted in SEQ ID NO. 1.

The proteins according to the invention are able to catalyze the conversion of a β -ionone structural element into a 3-hydroxy- β -ionone structural element, such as the conversion of β -carotene into zeaxanthin, β -carotene into β -cryptoxanthin,

β -cryptoxanthin into zeaxanthin, echinenone into 3'-hydroxyechinenone, 3-hydroxyechinenone into adonixanthin (4-ketozeaxanthin), α -carotene into α -cryptoxanthin or other chemical compounds which have up to 40 C atoms and contain a β -ionone ring into the corresponding 3-hydroxy- β -ionone compounds or the conversion of a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone structural element, such as the conversion of canthaxanthin into astaxanthin, canthaxanthin into phoenicoxanthin (adonirubin), phoenicoxanthin (adonirubin) into astaxanthin, echinenone into 3-hydroxyechinenone, 3'-hydroxyechinenone into adonixanthin (4-ketozeaxanthin) or other chemical compounds which have up to 40 C atoms and contain a 4-keto- β -ionone ring into the corresponding 3-hydroxy-4-keto- β -ionone compounds.

15

For illustration of the xanthophylls derived from β -carotene, reference is made to the biosynthesis scheme in Misawa et al., J. Biotechnol. 1998, 59, page 174 (top). The biosynthesis of lutein takes place starting from α -carotene via α -cryptoxanthin.

20 The proteins which comprise a sequence which is derived from amino acid sequence SEQ ID NO. 2 by substitution, insertion or deletion of amino acids and which have an homology of at least 50% at the amino acid level with the sequence SEQ ID NO.2 have an enzymatic activity for the conversion of β -carotene into zeaxanthin or canthaxanthin into astaxanthin, preferably in an activity comparable to that of the protein comprising the sequence SEQ ID NO. 2.

Substitution means replacement of one or more amino acids by one or more amino acids. The replacements are preferably those called conservative, in which the replaced amino acid has a similar property to the original amino acid, for example replacement of Glu by Asp, Gln via Asn, Val by Ile, Leu by Ile, Ser by Thr.

35 Deletion is the replacement of an amino acid by a direct linkage. Preferred positions for deletions are the termini of the polypeptide and the linkages between the individual protein domains.

40 Insertions are introductions of amino acids into the polypeptide chain, there formally being replacement of a direct linkage by one or more amino acids.

The homology between two proteins means identity of the amino acids over the entire length of each protein, which is calculated by comparison with the aid of the computer program GAP (UWGCG, University of Wisconsin, Genetic Computer Group, program

algorithm of Needleman and Wunsch, J. Mol. Biol. 1970, 48, 443-453), setting the following parameters:

Gap Weight:	12
5 Length Weight:	4
Average Match:	2.912
Average Mismatch:	-2.003

A protein which has a homology of at least 50% at the amino acid
10 level with the sequence of SEQ ID NO.2 accordingly means a protein which, in comparison of its sequence with the sequence SEQ ID NO.2 using the above program algorithm with the above set of parameters, has an identity of at least 50%, preferably 60%, particularly preferably 70%.

15 The carotene hydroxylase according to the invention has a homology with the known prokaryotic β -carotene hydroxylases of 29.9% (Flavobacterium), 36.8% (Erwinia uredovora), 38.5% (Erwinia herbicola), 35.0% (Alcaligenes) and 35.6% (Agrobacterium
20 aurantiacum), and a homology of 41.2% with the known eukaryotic β -carotene hydroxylase from Arabidopsis thaliana.

A preferred protein has an enzymatic activity for the conversion of β -carotene into zeaxanthin and an enzymatic activity for the
25 conversion of canthaxanthin into astaxanthin and comprises the amino acid sequence SEQ ID NO. 2 or a protein which is derived from this sequence by substitution, insertion or deletion of amino acids and which has a homology of at least 50% at the amino acid level with the sequence of SEQ ID NO.2.

30 These preferred proteins are able to catalyze the conversion of a β -ionone structural element into a 3-hydroxy- β -ionone structural element, such as the conversion of β -carotene into zeaxanthin, β -carotene into β -cryptoxanthin, β -cryptoxanthin into zeaxanthin,
35 echinenone into 3'-hydroxyechinenone, 3-hydroxyechinenone into adonixanthin (4-ketozeaxanthin), α -carotene into α -cryptoxanthin or other chemical compounds which have up to 40 C atoms and contain a β -ionone ring into the corresponding 3-hydroxy- β -ionone compounds and the conversion of a 4-keto- β -ionone structural
40 element into a 3-hydroxy-4-keto- β -ionone structural element, such as the conversion of canthaxanthin into astaxanthin, canthaxanthin into phoenicoxanthin (adonirubin), phoenicoxanthin (adonirubin) into astaxanthin, echinenone into
3-hydroxyechinenone, 3'-hydroxyechinenone into adonixanthin
45 (4-ketozeaxanthin) or other chemical compounds which have up to

40 C atoms and contain a 4-keto- β -ionone ring into the corresponding 3-hydroxy-4-keto- β -ionone compounds.

- A particularly preferred protein is the eucaryotic carotene hydroxylase from the green alga *Haematococcus pluvialis* Flotow NIES-144 having the sequence SEQ ID NO. 2. This particularly preferred protein is able to catalyze the conversion of a β -ionone structural element into a 3-hydroxy- β -ionone structural element, such as the conversion of β -carotene into zeaxanthin, β -carotene into β -cryptoxanthin, β -cryptoxanthin into zeaxanthin, echinenone into 3'-hydroxyechinenone, 3-hydroxyechinenone into adonixanthin (4-ketozeaxanthin), α -carotene into α -cryptoxanthin or other chemical compounds which have up to 40 C atoms and contain a β -ionone ring into the corresponding 3-hydroxy- β -ionone compounds and the conversion of a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone structural element, such as the conversion of canthaxanthin into astaxanthin, canthaxanthin into phoenicoxanthin (adonirubin), phoenicoxanthin (adonirubin) into astaxanthin, echinenone into 3-hydroxyechinenone, 3'-hydroxyechinenone into adonixanthin (4-ketozeaxanthin) or other chemical compounds which have up to 40 C atoms and contain a 4-keto- β -ionone ring into the corresponding 3-hydroxy-4-keto- β -ionone compounds.
- 25 The carotene hydroxylases can be prepared, as described hereinafter, by gene expression of the appropriate nucleic acids which encode these proteins from natural or genetically manipulated organisms.
- 30 The invention further relates to nucleic acids, referred to as carotene hydroxylase genes hereinafter, which encode the proteins according to the invention described above. Suitable nucleic acid sequences can be obtained by back-translation of the polypeptide sequence in accordance with the genetic code. The codons preferably used for this purpose are those frequently used in accordance with the organism-specific codon usage. The codon usage can easily be found by means of computer analyses of other, known genes in the relevant organism.
- If, for example, the protein is to be expressed in a plant, it is often advantageous to use the codon usage of the plant in the back-translation.

- A preferred nucleic acid has the sequence SEQ ID NO. 1. This nucleic acid is a eucaryotic cDNA from the green alga *Haematococcus pluvialis* Flotow NIES-144, which encodes the carotene hydroxylase of sequence SEQ ID NO. 2. Since the reading frame of the cDNA is open toward the 5' end, SEQ ID NO. 1

possibly does not represent the complete sequence of the cDNA. Expression of this cDNA leads to a functional protein. Any partial sequence missing out the 5' end can be made up in a manner known per se by analyzing overlapping cDNA fragments of 5 the cDNA library from *Haematococcus pluvialis* Flotow NIES-144.

It is known that the green alga *Haematococcus pluvialis* produces large amounts of astaxanthin under unfavorable environmental conditions such as a phosphate or nitrogen deficit or where the 10 light intensity is high, to protect from photooxidative stress (Kobayashi et al., Appl. Environ. Microbiol. 1993, 59, 867-873; Boussiba et al., Methods Enzymol 1992, 213, 386-391). This process is normally accompanied by a morphological change in which the vegetative cells of the green algae develop into cyst 15 cells.

Addition of sodium acetate and FeSO_4 and an increase in the light intensity induced astaxanthin biosynthesis and cyst cell formation in a suspension culture of *Haematococcus pluvialis* Flotow 20 NIES-144. The RNA was isolated from *Haematococcus pluvialis* in this stage to construct a cDNA library. The cDNA having the sequence SEQ ID NO 1 was isolated from this cDNA library.

All the aforementioned carotene hydroxylase genes can be prepared 25 in a manner known per se by chemical synthesis from the nucleotide building blocks such as, for example, by fragment condensation of individual overlapping, complementary nucleic acid building blocks of the double helix. Chemical synthesis of oligonucleotides is possible, for example, in a known manner by 30 the phosphoramidite method (Voet, Voet, 2nd edition, Wiley Press New York, pages 896-897). Addition of synthetic oligonucleotides and filling in gaps using the Klenow fragment of DNA polymerase and ligation reactions, and general cloning methods, are described in Sambrook et al. (1989), Molecular cloning: A 35 laboratory manual, Cold Spring Harbor Laboratory Press. The invention further relates to nucleic acid constructs comprising one of the carotene hydroxylase genes according to the invention described above, which are linked to one or more regulatory signals to increased gene expression. 40 These regulatory sequences are, for example, sequences to which inducers or repressors bind and thus regulate the expression of the nucleic acid. In addition to these new regulatory sequences or in place of these sequences it is possible for the natural regulation of these sequences still to be present in front of the 45 actual structural genes and, where appropriate, have been genetically modified so that the natural regulation has been switched off and expression of the genes has been increased.

However, the nucleic acid construct may also have a simpler structure, that is to say no additional regulatory signals are inserted in front of the aforementioned carotene hydroxylase genes, and the natural promoter with its regulation is not
5 deleted. Instead, the natural regulatory sequence is mutated so that regulation no longer takes place, and gene expression is increased. These modified promoters may also be placed alone in front of the natural genes to increase the activity. The nucleic acid construct may additionally advantageously contain one or
10 more so-called enhancer sequences functionally linked to the promoter, which make increased expression of the nucleic acid sequence possible. It is also possible to insert additional advantageous sequences at the 3' end of the DNA sequences, such as further regulatory elements or terminators. The aforementioned
15 carotene hydroxylase genes may be present in one or more copies in the gene construct.

Advantageous regulatory sequences for the nucleic acid constructs according to the invention, for the process described below for
20 preparing xanthophylls and for the genetically modified organisms described below are present, for example, in promoters such as cos, tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacI^q, T7, T5, T3, gal, trc, ara, SP6, λ -P_R or in the λ -P_L promoter, which are advantageously used in Gram-negative bacteria.

25 Further advantageous regulatory sequences are present, for example, in the Gram-positive promoters amy and SPO2, in the yeast or fungal promoters ADC1, MF α , AC, P-60, CYC1, GAPDH, TEF, rp28, ADH or in the plant promoters CaMV/35S [Franck et al., Cell 21(1980) 285-294], PRP1 [Ward et al., Plant.Mol. Biol.22(1993)],
30 SSU, OCS, leb4, usp, STLS1, B33, nos or in the ubiquitin or phaseolin promoter.

Particularly advantageous are those plant promoters which ensure specific expression in tissues or plant parts in which the
35 biosynthesis of carotenoids or precursors thereof takes place or in which the products advantageously accumulate.

Particular mention should be made of promoters for the whole plant on the basis of constitutive expression, such as, for
40 example, the CaMV promoter, the OCS promoter from Agrobacterium (octopine synthase), the NOS promoter from Agrobacterium (nopaline synthase), the ubiquitin promoter, promoters of vacuolar ATPase subunits or the promoter of a proline-rich protein from wheat (WO 9113991),
45

seed-specific promoters such as, for example, the phaseolin promoter and the USP promoter from *Vicia faba*, the promoter of the legumin gene from *Vicia* (leb4) or the Bce4 gene promoter from *Brassica* (WO 9113980),

5

specific promoters for green tissues such as, for example, the SSU promoter (small subunit) of rubisco (ribulose-1,5-bisphosphate carboxylase) or the STLS1 promoter (*Solanum tuberosum*, light harvesting system 1 from potato),

10

mesophyll-specific promoters such as, for example, the FBPase promoter of the cytosolic FBPase from potato (WO9705900),

15

specific promoters for tubers, storage roots or roots, such as, for example, the patatin promoter class I (B33), the promoter of the cathepsin D inhibitor from potato, the promoter of starch synthase (GBSS1) or the sporamin promoter,

20

fruit-specific promoters such as, for example, the fruit-specific promoter from tomato (EP409625),

fruit ripening-specific promoters such as, for example, the fruit ripening-specific promoter from tomato (WO 9421794),

25

flower-specific promoters such as, for example, the phytoene synthase promoter (WO9216635) or the promoter of the P-rr gene (WO9822593),

30

specific plastid or chromoplast promoters such as, for example, the RNA polymerase promoter (WO9706250) or

35

pathogen- or chemically inducible promoters such as, for example, the PRP1 promoter, a benzenesulfonamide-inducible (EP 388186), a tetracycline-inducible (Gatz et al., (1992) *Plant J.* 2,397-404), an abscisic acid-inducible (EP335528) or an ethanol- or cyclohexanone-inducible (WO9321334) promoter.

40

It is also possible and advantageous to use the promoter of phosphoribosyl-pyrophosphate amidotransferase from *Glycine max* (see also Genbank accession number U87999) or another node-specific promoter as in EP 249676.

It is possible in principle to use all natural promoters with their regulatory sequences like those mentioned above for the process according to the invention. It is additionally possible and advantageous to use synthetic promoters.

45

The nucleic acid construct may also contain other genes which are to be introduced into the organisms. These genes may be under separate regulation or under the same regulatory region as the above-described carotene hydroxylase genes. Examples of these
5 genes are other biosynthetic genes of carotenoid biosynthesis which make increased synthesis possible. Particular mention may be made of other genes of carotenoid biosynthesis which are biochemically limiting, such as the genes encoding phytoene synthase, phytoene desaturase, isopentyl-pyrophosphate isomerase
10 or β -cyclase.

Genes encoding an isopentyl-pyrophosphate isomerase are known, for example, from the organisms *Phaffia rhodozyma* and *Haematococcus pluvialis* (EP 769 551) or *Arabidopsis thaliana* and
15 *Tagetes* (Marigold) (WO 9736998).

Genes encoding a phytoene synthase are known, for example, from the organisms *Erwinia uredovora* (EP 393690), *Erwinia herbicola* (WO 9113078), tomato (WO 9109128), melon (WO 9602650),
20 *Flavobacterium* (EP 747483) or *Nicotiana* (US 5705624).

Genes encoding a phytoene desaturase are known, for example, from the organisms *Erwinia uredovora* (EP 393690), *Erwinia herbicola* (WO 9113078), *Nicotiana* (US 5539093) or *Flavobacterium* (EP
25 747483).

Genes encoding a β -cyclase are known, for example, from the organisms *Erwinia uredovora* (EP 393690), *Erwinia herbicola* (WO 9113078), *Flavobacterium* (EP 747483), tobacco and tomato (WO
30 9628014) or *Capsicum annuum* (WO 9636717).

The invention further relates to a process for preparing the genetically modified organisms described below, wherein the carotene hydroxylase genes according to the invention or the nucleic acid constructs according to the invention are introduced
35 into the genome of the initial organism. By initial organisms are meant the organisms before the genetic modification according to the invention.

The carotene hydroxylase genes according to the invention or the
40 nucleic acid constructs according to the invention can in principle be introduced by all methods known to the skilled worker into the initial organisms described below, which are genetically modified thereby.

They are advantageously introduced into the initial organisms or cells thereof by transformation, transfection, electroporation, using the so-called particle gun or by microinjection.

- 5 The skilled worker can find appropriate methods for microorganisms in the textbooks by Sambrook, J. et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, by F.M. Ausubel et al. (1994) Current protocols in molecular biology, John Wiley and Sons, by D.M. Glover et al.,
10 DNA Cloning Vol. 1, (1995), IRL Press (ISBN 019-963476-9), by Kaiser et al. (1994) Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press or Guthrie et al. Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, 1994, Academic Press.

- 15 Examples of advantageous methods which may be mentioned are those such as the introduction of the DNA by homologous or heterologous recombination, for example using the ura-3 gene, specifically the ura-3 gene from Ashbya, as described in the German Application DE
20 19801120.2, and/or by the REMI method (= "restriction enzyme mediated integration) which is described below.

- The REMI technique is based on the cotransformation of a linear DNA construct which has been cut at both ends with the same
25 restriction endonuclease, together with the restriction endonuclease which was used for this restriction of the DNA construct, into an organism. The restriction endonuclease then cuts the genomic DNA of the organism into which the DNA construct has been introduced together with the restriction enzyme. This
30 leads to an activation of the cell's own repair mechanisms. These repair mechanisms repair the strand breaks in the genomic DNA which have been caused by endonuclease, and during this also incorporate with a certain frequency the cotransformed DNA construct into the genome. Ordinarily, the restriction cleavage
35 sites are retained at both ends of the DNA during this.

- This technique was described by Bölker et al. (Mol Gen Genet, 248, 1995: 547-552) for the insertion mutagenesis of fungi. The method was used by Von Schiestl and Petes (Proc. Natl. Acad. Sci.
40 USA, 88, 1991: 7585-7589) to find out whether there is heterologous recombination in Saccharomyces. The method has been described by Brown et al. (Mol. Gen. Genet. 251, 1996: 75-80) for the stable transformation and regulated expression of an inducible reporter gene.

It is possible using the REMI method to position the nucleic acid fragments according to the invention or the aforementioned carotene hydroxylase genes according to the invention at transcriptionally active sites in the genome.

- 5 It is possible and advantageous to clone the nucleic acids together with at least one reporter gene into a DNA construct, which is introduced into the genome. This reporter gene ought to make detectability easy by a growth, fluorescence, chemo- or
10 bioluminescence assay or by a photometric measurement. Examples which may be mentioned of reporter genes are antibiotic resistance genes, hydrolase genes, fluorescent protein genes, bioluminescence genes, glucosidase genes, the luciferase gene, β -galactosidase gene, gfp gene, lipase gene, esterase gene,
15 peroxidase gene, β -lactamase gene, acetyl-, phospho- or adenylyltransferase gene. These genes make it possible easily to measure and quantify the transcription activity and thus the expression of the genes. This means that it is possible to identify sites in the genome which have a productivity differing
20 by up to a factor of 2.

- If it is intended to introduce a plurality of genes, such as, for example, further crt genes of carotenoid biosynthesis, into the organism, they can all be introduced together with a reporter
25 gene in a single vector, or each individual gene with a reporter gene can be introduced in one vector in each case, into the organism, it being possible to introduce the various vectors at the same time or successively. It is also possible to insert gene fragments coding for the respective activities in the REMI
30 techniques.

- Restriction enzymes suitable in principle for integrating the carotene hydroxylase genes or nucleic acid constructs according to the invention into the genome of initial organisms are all
35 known ones. Restriction enzymes which recognize only 4 base pairs as restriction cleavage site are less preferred because they cut too often in the genome or in the vector to be integrated, and preferred enzymes recognize 6, 7, 8 or more base pairs as cleavage site, such as BamHI, EcoRI, BglII, SphI, SpeI, XbaI,
40 XhoI, NcoI, SalI, ClaI, KpnI, HindIII, SacI, PstI, BpnI, NotI, SrfI or SfiI, to mention only a few of the possible enzymes. It is advantageous if the enzymes used no longer have cleavage sites in the DNA to be introduced; this increases the efficiency of integration. Ordinarily, 5 to 500 U, preferably 10 to 250,
45 particularly preferably 10 to 100 U of the enzymes are used in the REMI mixture. The enzymes are advantageously employed in an aqueous solution which contains substances for osmotic

stabilization, such as sugars such as sucrose, trehalose or glucose, polyols such as glycerol or polyethylene glycol, a buffer with an advantageous buffering in the range of pH 5 to 9, preferably 6 to 8, particularly preferably 7 to 8, such as tris, 5 MOPS, HEPES, MES or PIPES and/or substances to stabilize the nucleic acids, such as inorganic or organic salts of Mg, Cu, Co, Fe, Mn or Mo. It is also possible where appropriate for the substances to be present, such as EDTA, EDDA, DTT, β -mercaptoethanol or nuclease inhibitors. However, it is also 10 possible to carry out the REMI technique without these additions.

The process is carried out at a temperature in the range from 5 to 80°C, preferably from 10 to 60°C, particularly preferably from 20 to 40°C. Other known methods for destabilizing cell membranes 15 are suitable for the process, such as, for example, electroporation, fusion with loaded vesicles or destabilization with various alkali metal or alkaline earth metal salts such as lithium, rubidium or calcium salts, with lithium salts being preferred.

20 The nucleic acids can be used for the reaction according to the invention directly after isolation or after purification.

The introduction of the carotene hydroxylase genes according to 25 the invention or the nucleic acid constructs according to the invention into plants can in principle take place by all methods known to the skilled worker.

The transfer of foreign genes into the genome of a plant is 30 referred to as transformation. In this case, the described methods for the transformation and regeneration of plants from plant tissues or plant cells for transient or stable transformation are utilized.

35 Suitable methods are protoplast transformation by polyethylene glycol-induced DNA uptake, the use of a gene gun, electroporation, incubation of dry embryos in DNA-containing solution, microinjection and *Agrobacterium*-mediated gene transfer. The methods mentioned are described, for example, in B. 40 Jenes et al., *Techniques for Gene Transfer*, in: *Transgenic Plants*, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press (1993) 128-143, and in Potrykus *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42 (1991) 205-225.

45 The construct to be expressed is preferably cloned into a vector which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., *Nucl. Acids Res.* 12 (1984) 8711).

Transformation with *Agrobacterium tumefaciens* is described, for example, by Höfgen and Willmitzer in Nucl. Acids Res. (1988) 16, 9877.

- 5 *Agrobacteria* transformed with an expression vector according to the invention can likewise be used in a known manner for transforming plants, in particular tagetes, sunflower, arabidopsis, tobacco, red pepper, soybean, tomato, aubergine, paprika, carrot, potato, corn, lettuce and brassica species,
10 oats, rye, wheat, triticale, millet, rice, alfalfa, flax, brassicaceae such as, for example, oilseed rape or canola, sugar beet, sugar cane or wood plants such as, for example, aspen or yew, for example by bathing wounded leaves or pieces of leaves in a solution of *agrobacteria* and then cultivating in suitable
15 media.

The genetically modified plant cells can be regenerated by all methods known to the skilled worker. Appropriate methods can be found in the abovementioned publications by S.D. Kung and R. Wu,
20 Potrykus or Höfgen and Willmitzer.

There is a large number of possibilities for increasing the enzymic activity of the carotene hydroxylase gene products in the cell.

- 25 One possibility is to modify the endogenous carotene hydroxylase genes so that they code for enzymes having a carotene hydroxylase activity which is higher than the initial enzymes. Another increase in the enzymic activity can be achieved, for example, by
30 modifying the catalytic sites to increase substrate conversion or by aborting the effect of enzyme inhibitors, that is to say they have an increased specific activity or their activity is not inhibited. The enzymic activity can also be increased in a further advantageous embodiment by increasing enzyme synthesis in
35 the cell, for example by eliminating factors which repress enzyme synthesis or by increasing the activity of factors or regulatory elements which promote enhanced synthesis, or preferably by introducing further gene copies. This measure increases the total activity of the gene products in the cell without altering the
40 specific activity. It is also possible to use a combination of these methods, that is to say increasing the specific activity plus increasing the total activity. These modifications can in principle be introduced by all methods known to the skilled worker into the nucleic acid sequences of the genes, regulatory
45 elements or promoters thereof. It is possible for this purpose to subject the sequences, for example, to a mutagenesis such as a site directed mutagenesis as described in D.M. Glover et al., DNA

Cloning Vol.1, (1995), IRL Press (ISBN 019-963476-9), chapter 6, page 193 et seq.

Spee et al. (Nucleic Acids Research, Vol. 21, No. 3, 1993: 777 - 778) describes a PCR method using dITP for random mutagenesis.

The use of an in vitro recombination technique for molecular evolution is described by Stemmer (Proc. Natl. Acad. Sci. USA, Vol. 91, 1994: 10747 - 10751).

10

Moore et al. (Nature Biotechnology Vol. 14, 1996: 458 - 467) describe combination of the PCR and recombination methods.

The modified nucleic acid sequences are subsequently returned to the organisms via vectors.

15

For increasing enzymic activities it is also possible to place modified promoter regions in front of the natural genes so that expression of the genes is increased and thus the activity is eventually raised. Sequences can also be introduced at the 3' end, for example to increase the stability of the mRNA and thus make increased translation possible. This likewise leads to higher enzymic activity.

20

In a further preferred embodiment, the carotene hydroxylase genes or nucleic acid constructs according to the invention are, for expression in one of the organisms described below, inserted into a vector such as, for example, a plasmid, a phage or other DNA which makes optimal expression of the genes possible in the prokaryotic or eukaryotic organisms. This vector may contain the ATG start codon for expression of the protein according to the invention.

25

30

Examples of suitable plasmids in E. coli are pLG338, pACYC184, pBR322, pUC18, pUC19, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III¹¹³-B1, λ gt11, pBdCI, the pET vector series (Novagen and Stratagene), pMAL or the pQE vector series (Qiagen), in fungi pALS1, pIL2 or pBB116, in yeasts 2 μ , pAG-1, YEpl6, YEpl3 or pEMBLye23 or in plants pLGV23, pGHLac⁺, pBIN19, pAK2004, pDH51, or derivatives of the aforementioned plasmids.

35

40

Said plasmids represent a small selection of the possible plasmids. Further plasmids are well known to the skilled worker and can be found, for example, in the book Cloning Vectors (Eds. Pouwels P. H. et al. Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). Suitable plant vectors are described inter

45

alia in "Methods in Plant Molecular Biology and Biotechnology" (CRC Press), chapters 6/7, pp.71-119.

It is advantageous for expression of the other genes present if the nucleic acid fragment also contains in addition 3' and/or 5' terminal regulatory sequences to increase expression, which are selected for optimal expression depending on the selected host organism and gene or genes.

10 These regulatory sequences are intended to make specific expression of the genes and protein expression possible. This may mean, for example, depending on the host organism that the gene is expressed and/or overexpressed only after induction, or that it is immediately expressed and/or overexpressed.

15 The regulatory sequences or factors may moreover preferably have a beneficial effect on expression of the introduced genes, and thus increase it. Enhancement of the regulatory elements is thus advantageously possible at the level of transcription by using
20 strong transcription signals such as promoters and/or enhancers. However, it is also possible to enhance translation by, for example, improving the stability of the mRNA.

In a further embodiment of the vector, the nucleic acid construct
25 according to the invention may also advantageously be introduced in the form of a linear DNA into the host organisms and be integrated into the genome of the host organism by heterologous or homologous recombination. This linear DNA may consist of a linearized plasmid or only of the nucleic acid fragment as
30 vector.

It is also possible to use as vector any plasmid which undergoes autonomous replication in the cell, but also, as described above, a linear DNA fragment which integrates into the genome of the
35 host. This integration can take place by heterologous or homologous recombination, but preferably, as mentioned, by homologous recombination (Steiner et al., Genetics, Vol. 140, 1995: 973-987). It is moreover possible for the aforementioned carotene hydroxylase genes to be present singly in the genome at
40 various sites or on various vectors or to be present together in the genome or on one vector.

The invention further relates to the use of the carotene hydroxylase genes according to the invention for producing
45 genetically modified organisms.

The invention further relates to a correspondingly genetically modified organism, with the expression of the carotene hydroxylase genes according to the invention being increased by comparison with a wild type in the case where
5 the initial organism contains a carotene hydroxylase gene according to the invention,
or being caused in the case where the initial organism does not contain a carotene hydroxylase gene according to the invention, by the genetic modification.

10

A genetically modified organism means an organism in which the carotene hydroxylase genes or nucleic acid constructs according to the invention have been inserted, preferably by one of the methods described above.

15

The genetically modified organism contains at least one carotene hydroxylase gene according to the invention or at least one nucleic acid construct according to the invention. Depending on the initial organism, the nucleic acid may be present inside or
20 outside the chromosome.

Carotenoid metabolism in the genetically modified organisms is preferably altered by comparison with the wild type.

25 Suitable genetically modified organisms are in principle all organisms able to synthesize xanthophylls.

Preferred initial organisms are those naturally able to synthesize xanthophylls. However, initial organisms able to
30 synthesize xanthophylls because of the introduction of carotenoid biosynthesis genes are also suitable.

Initial organisms mean prokaryotic or eukaryotic organisms such as, for example, microorganisms or plants. Preferred microorganisms are bacteria, yeasts, algae or fungi.

35

Bacteria which can be used are both bacteria which are able, because of the introduction of carotenoid biosynthesis genes of a carotenoid-producing organism to synthesize xanthophylls, such as, for example, bacteria of the genus *Escherichia*, which
40 contain, for example, crt genes from *Erwinia*, and bacteria which are intrinsically able to synthesize xanthophylls, such as, for example, bacteria of the genus *Erwinia*, *Agrobacterium*, *Flavobacterium*, *Alcaligenes* or *Cyanobacteria* of the genus *Synechocystis*. Preferred bacteria are *Escherichia coli*, *Erwinia*
45 *herbicola*, *Erwinia uredovora*, *Agrobacterium aurantiacum*,

Alcaligenes sp. PC-1, *Flavobacterium* sp. strain R1534 or the *Cyanobacterium* *Synechocystis* sp. PCC6803.

Preferred uses are *Candida*, *Saccharomyces*, *Hansenula* or *Pichia*.

5

Preferred fungi are *Aspergillus*, *Trichoderma*, *Ashbya*, *Neurospora*, *Fusarium* or other fungi described in Indian Chem Engr. Section B. Vol 37, No. 1,2 (1995) on page 15, Table 6.

10 Preferred algae are green algae such as, for example, algae of the genus *Haematococcus*, *Phaedactylum tricornatum*, *Volvox* or *Dunaliella*. Particular preferred algae are *Haematococcus pluvialis* or *Dunaliella bardawil*.

15 In a preferred embodiment, plants are used as initial organisms and, accordingly, also as genetically modified organisms. Examples of preferred plants are tagetes, sunflower, arabidopsis, tobacco, red pepper, soybean, tomato, aubergine, paprika, carrot, potato, corn, lettuce and brassica species, oats, rye, wheat, 20 triticale, millet, rice, alfalfa, flax, brassicaceae such as, for example, oilseed rape or canola, sugar beet, sugar cane or woody plants such as, for example aspen or yew.

Particular preference is given to *Arabidopsis thaliana*, *Tagetes* 25 *erecta*, oilseed rape, canola, potatoes, and oilseeds and typical carotenoid producers such as soybean, sunflower, paprika, carrot, pepper or corn.

The invention further relates to a process for the preparation of 30 xanthophyll derivatives, which comprises converting a β -ionone structural element into a 3-hydroxy- β -ionone structural element and/or a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone structural element in the presence of the protein according to the invention.

35

Xanthophyll derivatives mean xanthophylls, preferably xanthophylls containing at least one hydroxyl group, such as, for example, zeaxanthin, β -cryptoxanthin, 3'-hydroxyechinenone, 3-hydroxyechinenone, adonixanthin (4-ketozeaxanthin), 40 astaxanthin, phoenicoxanthin (adonirubin), α -cryptoxanthin or lutein or derivatives thereof having up to 40 C atoms and containing at least one 3-hydroxy- β -ionone or at least one 3-hydroxy-4-keto- β -ionone structural element in the molecule, such as, for example, 3-hydroxy-6-vinyl- β -ionone, 45 3-hydroxy-4-keto-6-vinyl- β -ionone, 3-hydroxyretinol, 3-hydroxy-4-ketoretinol, 3-hydroxyretinal,

3-hydroxy-4-ketoretinal, 3-hydroxyretinoic acid or 3-hydroxy-4-ketoretinoic acid.

Preferred xanthophyll derivatives are zeaxanthin, lutein and 5 astaxanthin.

In the process according to the invention there is conversion in the presence of the proteins according to the invention of a β -ionone structural element into a 3-hydroxy- β -ionone structural
10 element, such as β -carotene into zeaxanthin, β -carotene into β -cryptoxanthin, β -cryptoxanthin into zeaxanthin, echinenon into 3'-hydroxyechinenone, 3-hydroxyechinenone into adonixanthin (4-ketozeaxanthin), α -carotene into α -cryptoxanthin or a chemical compound having up to 40 C atoms and containing a β -ionone ring
15 into the corresponding 3-hydroxy- β -ionone compound or a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone structural element, such as canthaxanthin into astaxanthin, canthaxanthin in phoenicoxanthin (adonirubin), phoenicoxanthin (adonirubin) into astaxanthin,
20 echinenone into 3-hydroxyechinenone, 3'-hydroxyechinenone into adonixanthin (4-ketozeaxanthin) or a chemical compound having up to 40 C atoms and containing a 4-keto- β -ionone ring into the corresponding 3-hydroxy-4-keto- β -ionone compound.

25 In a preferred embodiment of the process, an aforementioned genetically modified organism according to the invention is cultivated, the latter organism is harvested, and subsequently the xanthophyll derivatives are isolated from the organism.

30 Cultivation of the genetically modified organism according to the invention takes place in a manner known per se, such as cultivation of the appropriate wild type, for example in the case of microorganisms in a suitable medium such as, for example, on agar plates or in suspension culture, or in the case of plants in
35 soil or appropriately suitable nutrient media.

By harvesting is meant in the case of microorganisms the isolation of the microorganisms, and in the case of plants the cutting off of the plant or, where appropriate, particular plant parts containing the xanthophyll derivatives.

40 The xanthophyll derivatives are isolated in a manner known per se, for example by disruption of the organism cells, extraction of the xanthophyll derivatives and subsequent purification of the xanthophyll derivatives by chemical or physical separation methods such as extraction or chromatography.

The invention further relates to the use of the carotene hydroxylase according to the invention or of the carotene hydroxylase genes according to the invention for preparing xanthophyll derivatives.

5

The following examples illustrate the invention

Example 1

Incorporation of the carotene hydroxylase gene into a β -carotene
10 producing *E. coli*, fermentation of the transgenic organism and
isolation of the xanthophylls

General methods

15 Isolation of the carotenoids

To isolate the carotenoids (carotenes and xanthophylls), the *E. coli* cells were collected by centrifugation, and the resulting cell material was freeze dried for 24 h. The freeze-dried cells
20 were resuspended in acetone and extracted twice with acetone at 55°C for 15 min. The combined extracts were washed with a diethyl ether/petroleum ether (boiling point 35-80°C) mixture (1:9, v/v) in a separating funnel and concentrated to dryness under nitrogen in a rotary evaporator.

25

HPLC analysis

The extracts were fractionated using a Nucleosil 100-5 C18 column (Macherey-Nagel) with an eluent flow rate of 1.5 ml/min.
30 The eluent used to fractionate the β -carotene and the hydroxylated xanthophylls in Example 1 was an acetonitrile/methanol/2-propanol mixture (85:10:5, v/v/v; flow).

To fractionate the xanthophylls with keto groups in Example 2, an acetonitrile/methanol/H₂O mixture (50:44:6, v/v/v) was used as
35 eluent 1 for 22 min, and methanol was used as eluent 2. Detection took place directly using the Waters 994 diode array detector. A comparison standard for the HPLC analysis were β -carotene, astaxanthin and zeaxanthin purchased from Sigma or Roth.

40 1.1 Production of β -carotene-producing *E. coli*

The organism used comprised *E. coli* cells of the strain JM101. The plasmid pACCAR16DcrtX contains the bacterial carotenoid biosynthesis genes *crtE*, *crtB*, *crtI* and *crtY* from *Erwinia*
45 *uredovora* and results in the biosynthesis of β -carotene (N. Misawa et al., J. Bacteriol. 172 (1990) 6704-6712; Biochem. Biophys. Res. Commun. 209 (1995) 867-876)

- To prepare this plasmid, a 6.0 kb Asp718(KpnI)-EcoRI fragment of the carotenoid biosynthesis gene cluster from *Erwinia uredovora* (Plasmid pCAR16delB) was cloned into the EcoRI site of the plasmid pACY184 (R.E. Rose, Nucl. Acids Res. 16 (1988) 355). The
- 5 plasmid pCAR16delB contains a frameshift mutation in the β -carotene hydroxylase ORF (open reading frame) (Misawa et al., Biochem. Biophys. Res. Commun. 209 (1995) 867-876). It is therefore impossible for the resulting β -carotene to be hydroxylated to zeaxanthin.
- 10 Insertion of the plasmid pACCAR16DcrtX into *E. coli* and preparation and isolation of transformed *E. coli* cells took place in a manner known per se, as described in Sambrook et al., Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.
- 15
- 1.2 Construction of a *Haematococcus pluvialis* λ cDNA expression library and isolation of the plasmid containing the carotene hydroxylase gene
- 20 *Haematococcus pluvialis* Flotow NIES-144 originated from the National Institute for Environmental Studies (NIES), Tsukuba, Japan. The basal medium (pH 6.8) in the growth phase of *Haematococcus pluvialis* contained per liter 1.2 g of sodium acetate, 2.0 g of yeast extract, 0.4 g of L-asparagine, 0.2 g of
- 25 $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 0.01 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, and 0.02 g of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$. *Haematococcus pluvialis* was grown at 20°C with a dark/light cycle of 12 h light (20 $\mu\text{E}/\text{m}^2\text{s}$) and 12 h dark for 4 days. To induce astaxanthin biosynthesis and cyst cell formation, after 4 days sodium acetate and FeSO_4 were added to a concentration of
- 30 45 mM and 450 μM respectively. After the addition, the light conditions were changed to continuous light (125 $\mu\text{E}/\text{m}^2\text{s}$) as described in Kajiwara et al., Plant Mol. Biol. 29 (1995) 343-352. After induction of cyst cell formation for 8 h, the RNAs of *Haematococcus pluvialis* were isolated to construct a cDNA library
- 35 from the cyst cells.
- The poly(A)RNA was purified using oligo(dT)-cellulose (Biolabs). Synthesis of the cDNAs and construction of the λ ZAP expression library took place with the aid of the cDNA Synthesis and ZAP-cDNA Gigapack III Gold Cloning Kit (Stratagene). The first
- 40 strand of the cDNA was synthesized using MMLV reverse transcriptase and a poly(dT) primer containing an XhoI restriction enzyme recognition site in accordance with the Stratagene instructions for use. The second strand was synthesized correspondingly using DNA polymerase I. Blunt DNA
- 45 ends were produced by filling in with Pfu DNA polymerase, and EcoRI adapters were ligated thereto. The resulting cDNA fragments were then fractionated according to size and subsequently ligated

into the EcoRI - XhoI recognition site of the Uni-ZAP XR vector. After isolation and purification of the positive carotene hydroxylase plaque, the pBluescript phagemid containing the carotene hydroxylase cDNA was recovered by in vivo cutting out by use of the ExAssist helper phage and the SOLR E. coli strain, in accordance with the Stratagene instructions for use. The resulting plasmid contains in addition to the short adapter sequences at the 5'- (5'-AATTCGGCAG-3') and 3' (5'-TCGAG-3') ends according to DNA sequence analysis a 1608 bp-long cDNA fragment ligated into the EcoRI and XhoI restriction site of the multiple cloning site. This plasmid was used for insertion of the carotene hydroxylase gene into the β -carotene-producing E. coli cells as described under 1.1.

1.3. Incorporation of the carotene hydroxylase gene into β -carotene-producing E. coli cells, cultivation of the transformed cells and isolation of the xanthophylls zeaxanthin and β -cryptoxanthin

Transformation of the β -carotene-producing E. coli cells containing the plasmid pACCAR16DcrtX, described under 1.1., with the plasmid containing the carotene hydroxylase gene, described under 1.3., took place in a manner known per se as described in J. Sambrook et al., Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The transformed E. coli cells were cultured in LB medium at 28°C with addition of ampicillin (50 μ g/ml; for the plasmid containing the carotene hydroxylase gene) and chloramphenicol (30 μ g/ml; plasmid pACCAR16DcrtX) for 48 h.

The carotenoids were isolated as described under the general methods. Fig. 1 shows the HPLC diagram for the carotenoids extracted from

(A) E. coli cells containing the plasmid pACCAR16DcrtX from 1.1

and

(B) E. coli cells containing the plasmid pACCAR16DcrtX together with the carotene hydroxylase gene from 1.2.

The peak numbers in Fig. 1 mean

3 zeaxanthin
5 β -cryptoxanthin
6 β -carotene

5 As is evident from Fig. 1, the transformed *E. coli* cells produce, due to incorporation of the carotene hydroxylase gene according to the invention, the hydroxylated xanthophylls zeaxanthin and β -cryptoxanthin, whereas only β -carotene is produced without the transformation. The carotene hydroxylase according to the
10 invention is accordingly able to convert β -carotene into β -cryptoxanthin, β -cryptoxanthin into zeaxanthin or β -carotene into zeaxanthin.

Example 2

15 Incorporation of the carotene hydroxylase gene and the β -carotene ketolase gene (bkt) from *Haematococcus pluvialis* into β -carotene-producing *E. coli* cells, cultivation of the transformed cells and isolation of the xanthophylls astaxanthin, canthaxanthin, adonixanthin, zeaxanthin and β -cryptoxanthin

20

2.1 Preparation of a plasmid containing the β -carotene ketolase gene (bkt) from *Haematococcus pluvialis*

To prepare the plasmid pRKbkt1 (Kajiwara et al., Plant Mol. Biol.
25 29 (1995) 343-352) containing the β -carotene ketolase gene from *Haematococcus pluvialis*, a 1 kb product of a PvuII partial digest of the pUC19bkt plasmid was fractionated by agarose gel electrophoresis (Breitenbach et al., FEMS Microbiol. Lett. 140 (1996) 241-246). This DNA fragment contains the lacZ promoter
30 together with the ORF (open reading frame) of the ketolase and was subcloned into the plasmid pRK404 which had previously been digested with HindIII and treated with the Klenow enzyme.

The resulting plasmid pRKbkt1 was inserted once alone and once
35 together with the carotene hydroxylase gene plasmid described under 1.2 into the β -carotene-producing *E. coli* cells.

2.2 Incorporation of the plasmid pRKbkt1 into β -carotene-producing
40 *E. coli* cells, cultivation of the transformed cells and isolation of the xanthophyll canthaxanthin

Transformation of the β -carotene-producing *E. coli* cells containing the plasmid pACCAR16DcrtX, described under 1.1., with the plasmid pRKbkt1 containing the β -carotene ketolase gene (bkt),
45 described under 2.1., took place in a manner known per se as described in J. Sambrook et al., Molecular cloning: a laboratory

manual, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The transformed *E. coli* cells were cultured in analogy to the description under 1.3. in LB medium at 28 °C, but with the addition of chloramphenicol (30 µg/ml; plasmid pACCAR16DcrtX), tetracycline (10 µg/ml; plasmid pRKbkt1) and isopropyl β-D-thio-galactopyranoside (0.5 mM) for 48 h.

- 10 The carotenoids were isolated as described under the general methods.

Fig. 2 (A) shows HPLC-diagram of the carotenoids extracted from *E. coli* cells containing the plasmid pACCAR16DcrtX together with the plasmid pRKbkt1, containing the β-carotene ketolase gene (bkt).

Incorporation of the plasmid pRKbkt1 results in the transformed *E. coli* cells producing the xanthophyll with a keto group canthaxanthin.

20

2.3 Incorporation of the plasmid pRKbkt1 and the carotene hydroxylase gene into β-carotene-producing *E. coli* cells, cultivation of the transformed cells and isolation of the xanthophylls, astaxanthin, canthaxanthin, adonixanthin,

- 25 zeaxanthin and β-cryptoxanthin

Transformation of the β-carotene-producing *E. coli* cells containing the plasmid pACCAR16DcrtX, described under 1.1., with the plasmid pRKbkt1 containing the β-carotene ketolase gene (bkt), described under 2.1., with the plasmid containing the carotene hydroxylase gene, described under 1.3, took place in a manner known per se as described in J. Sambrook et al., Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

35

The transformed *E. coli* cells were cultured in analogy to the description under 1.3. in LB medium at 28°C, but with the addition of ampicillin (50 µg/ml; plasmid containing the carotene hydroxylase gene), chloramphenicol (30 µg/ml; plasmid pACCAR16DcrtX), tetracycline (10 µg/ml; plasmid pRKbkt1) and isopropyl β-D-thio-galactopyranoside (0.5 mM) for 48 h.

The carotenoids were isolated as described under the general methods. Fig. 2 shows the HPLC diagrams for the carotenoids extracted from

45

(A) *E. coli* cells containing the plasmid pACCAR16DcrtX from 1.1. together with the plasmid pRKbkt1 and

(B) *E. coli* cells containing the plasmid pACCAR16DcrtX together
5 with the plasmid pRKbkt1 and the carotene hydroxylase gene from 1.2.

Fig. 2 (C) shows astaxanthin as comparison standard.

10 The peak numbers in Fig. 2 mean

1	astaxanthin
2	adonixanthin
3	zeaxanthin
15 4	canthaxanthin
5	β -cryptoxanthin
6	β -carotene

As can be seen in Fig. 2, the transformed *E. coli* cells produce,
20 due to incorporation of the carotene hydroxylase gene according to the invention together with the plasmid pRKbkt1 containing the β -carotene ketolase gene (bkt), the hydroxylated and/or keto-containing xanthophylls astaxanthin, canthaxanthin, adonixanthin, zeaxanthin and β -cryptoxanthin, whereas only
25 canthaxanthin is produced without the carotene hydroxylase gene according to the invention.

In enzyme studies on the β -carotene ketolase from *Haematococcus pluvialis* it was shown that the enzyme mainly converts β -carotene
30 into canthaxanthin, whereas hydroxyl-containing xanthophylls such as zeaxanthin and β -cryptoxanthin are scarcely converted, and only to adonixanthin and not to astaxanthin (T. Lotan, J. Hirschberg, FEBS Lett. 364 (1995) 125-128; J. Breitenbach, N. Misawa, S. Kajiwara, G. Sandmann, FEMS Microbiol. Lett. 140 (1996) 241-246;
35 P.D. Fraser, H. Shimada, N. Misawa, Eur. J. Biochem. 252 (1998) 229-236).

The fact that the *E. coli* cells transformed in Example 2.3 are able to produce astaxanthin proves that the carotene hydroxylase
40 according to the invention is able to convert canthaxanthin via phoenicoxanthin (adonirubin) into astaxanthin.

The carotene hydroxylase according to the invention is accordingly able to catalyze the conversion of a β -ionone
45 structural element into a 3-hydroxy- β -ionone structural element, such as the conversion of β -carotene into zeaxanthin, β -carotene into β -cryptoxanthin, β -cryptoxanthin into zeaxanthin, echinenone

- into 3'-hydroxyechinenone, 3-hydroxyechinenone into adonixanthin (4-ketozeaxanthin), α -carotene into α -cryptoxanthin or other chemical compounds which have up to 40 C atoms and contain a β -ionone ring into the corresponding 3-hydroxy- β -ionone compounds
- 5 or the conversion of a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone structural element, such as the conversion of canthaxanthin into astaxanthin, canthaxanthin into phoenicoxanthin (adonirubin), phoenicoxanthin (adonirubin) into astaxanthin, echinenone into 3-hydroxyechinenone,
- 10 3'-hydroxyechinenone into adonixanthin (4-ketozeaxanthin) or other chemical compounds which have up to 40 C atoms and contain a 4-keto- β -ionone ring into the corresponding 3-hydroxy-4-keto- β -ionone compounds.

15

20

25

30

35

40

45

SEQUENCE LISTING

<110> BASF Aktiengesellschaft

<120> Carotene hydroxylase and process for preparing xanthophyll derivatives

<130> OZ 0050/49896

<140>

<141>

<160> 2

<170> PatentIn Vers. 2.0

<210> 1

<211> 1607

<212> DNA

<213> Haematococcus pluvialis

<220>

<221> CDS

<222> (3)..(968)

<400> 1

ct	aca	ttt	cac	aag	ccc	gtg	agc	ggt	gca	agc	gct	ctg	ccc	cac	atc	47
	Thr	Phe	His	Lys	Pro	Val	Ser	Gly	Ala	Ser	Ala	Leu	Pro	His	Ile	
	1				5				10				15			

ggc	cca	cct	ctt	cat	ctc	cat	cgg	tca	ttt	gct	gct	acc	acg	atg	ctg	95
Gly	Pro	Pro	Pro	His	Leu	His	Arg	Ser	Phe	Ala	Ala	Thr	Thr	Met	Leu	
				20				25					30			

tcg	aag	ctg	cag	tca	atc	agc	gtc	aag	gcc	cgc	cgc	gtt	gaa	cta	gcc	143
Ser	Lys	Leu	Gln	Ser	Ile	Ser	Val	Lys	Ala	Arg	Arg	Val	Glu	Leu	Ala	
		35					40					45				

cgc	gac	atc	acg	cgg	ccc	aaa	gtc	tgc	ctg	cat	gct	cag	cgg	tgc	tcg	191
Arg	Asp	Ile	Thr	Arg	Pro	Lys	Val	Cys	Leu	His	Ala	Gln	Arg	Cys	Ser	
		50					55					60				

tta	gtt	cgg	ctg	cga	gtg	gca	gca	cca	cag	aca	gag	gag	gcg	ctg	gga	239
Leu	Val	Arg	Leu	Arg	Val	Ala	Ala	Pro	Gln	Thr	Glu	Glu	Ala	Leu	Gly	
	65				70					75						

acc	gtg	cag	gct	gcc	ggc	gcg	ggc	gat	gag	cac	agc	gcc	gat	gta	gca	287
Thr	Val	Gln	Ala	Ala	Gly	Ala	Gly	Asp	Glu	His	Ser	Ala	Asp	Val	Ala	
	80				85				90				95			

ctc	cag	cag	ctt	gac	cgg	gct	atc	gca	gag	cgt	cgt	gcc	cgg	cgc	aaa	335
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Leu	Gln	Gln	Leu	Asp	Arg	Ala	Ile	Ala	Glu	Arg	Arg	Ala	Arg	Arg	Lys	
				100					105						110	
cgg	gag	cag	ctg	tca	tac	cag	gct	gcc	gcc	att	gca	gca	tca	att	ggc	383
Arg	Glu	Gln	Leu	Ser	Tyr	Gln	Ala	Ala	Ala	Ile	Ala	Ala	Ser	Ile	Gly	
			115					120					125			
gtg	tca	ggc	att	gcc	atc	ttc	gcc	acc	tac	ctg	aga	ttt	gcc	atg	cac	431
Val	Ser	Gly	Ile	Ala	Ile	Phe	Ala	Thr	Tyr	Leu	Arg	Phe	Ala	Met	His	
		130					135					140				
atg	acc	gtg	ggc	ggc	gca	gtg	cca	tgg	ggt	gaa	gtg	gct	ggc	act	ctc	479
Met	Thr	Val	Gly	Gly	Ala	Val	Pro	Trp	Gly	Glu	Val	Ala	Gly	Thr	Leu	
	145					150				155						
ctc	ttg	gtg	gtt	ggt	ggc	gcg	ctc	ggc	atg	gag	atg	tat	gcc	cgc	tat	527
Leu	Leu	Val	Val	Gly	Gly	Ala	Leu	Gly	Met	Glu	Met	Tyr	Ala	Arg	Tyr	
160				165					170						175	
gca	cac	aaa	gcc	atc	tgg	cat	gag	tcg	cct	ctg	ggc	tgg	ctg	ctg	cac	575
Ala	His	Lys	Ala	Ile	Trp	His	Glu	Ser	Pro	Leu	Gly	Trp	Leu	Leu	His	
				180				185					190			
aag	agc	cac	cac	aca	cct	cgc	act	gga	ccc	ttt	gaa	gcc	aac	gac	ttg	623
Lys	Ser	His	His	Thr	Pro	Arg	Thr	Gly	Pro	Phe	Glu	Ala	Asn	Asp	Leu	
			195					200					205			
ttt	gca	atc	atc	aat	gga	ctg	ccc	gcc	atg	ctc	ctg	tgt	acc	ttt	ggc	671
Phe	Ala	Ile	Ile	Asn	Gly	Leu	Pro	Ala	Met	Leu	Leu	Cys	Thr	Phe	Gly	
	210					215						220				
ttc	tgg	ctg	ccc	aac	gtc	ctg	ggg	gcg	gcc	tgc	ttt	gga	gcg	ggg	ctg	719
Phe	Trp	Leu	Pro	Asn	Val	Leu	Gly	Ala	Ala	Cys	Phe	Gly	Ala	Gly	Leu	
	225					230				235						
ggc	atc	acg	cta	tac	ggc	atg	gca	tat	atg	ttt	gta	cac	gat	ggc	ctg	767
Gly	Ile	Thr	Leu	Tyr	Gly	Met	Ala	Tyr	Met	Phe	Val	His	Asp	Gly	Leu	
240				245				250						255		
gtg	cac	agg	cgc	ttt	ccc	acc	ggg	ccc	atc	gct	ggc	ctg	ccc	tac	atg	815
Val	His	Arg	Arg	Phe	Pro	Thr	Gly	Pro	Ile	Ala	Gly	Leu	Pro	Tyr	Met	
				260				265					270			
aag	cgc	ctg	aca	gtg	gcc	cac	cag	cta	cac	cac	agc	ggc	aag	tac	ggt	863
Lys	Arg	Leu	Thr	Val	Ala	His	Gln	Leu	His	His	Ser	Gly	Lys	Tyr	Gly	
			275				280						285			
ggc	gcg	ccc	tgg	ggt	atg	ttc	ttg	ggt	cca	cag	gag	ctg	cag	cac	att	911
Gly	Ala	Pro	Trp	Gly	Met	Phe	Leu	Gly	Pro	Gln	Glu	Leu	Gln	His	Ile	
	290					295						300				

cca ggt gcg gcg gag gag gtg gag cga ctg gtc ctg gaa ctg gac tgg 959
 Pro Gly Ala Ala Glu Glu Val Glu Arg Leu Val Leu Glu Leu Asp Trp
 305 310 315

tcc aag cgg tagggtgcgg aaccaggcac gctggtttca cacctcatgc 1008
 Ser Lys Arg
 320

ctgtgataag gtgtggctag agcgatgcgt gtgagacggg tatgtcacgg tcgactggtc 1068
 tgatggccaa tggcatcggc catgtctggt catcacgggc tggttgcctg ggtgaagggtg 1128
 atgcacatca tcatgtgcgg ttggaggggc tggcacagtg tgggctgaac tggagcagtt 1188
 gtccaggctg gcgttgaatc agtgagggtt tgtgattggc ggttgtgaag caatgactcc 1248
 gcccatattc tatttgtggg agctgagatg atggcatgct tgggatgtgc atggatcatg 1308
 gtagtgcagc aaactatatt cacctagggc tgttggtagg atcagggtgag gccttgacaca 1368
 ttgcatgatg tactcgtcat ggtgtgttgg tgagaggatg gatgtggatg gatgtgtatt 1428
 ctcagacgta gaccttgact ggaggcttga tcgagagagt gggccgtatt ctttgagagg 1488
 ggaggctcgt gccagaaatg gtgagtggat gactgtgacg ctgtacattg caggcagggtg 1548
 agatgcactg tctcgattgt aaaatacatt cagatgcaaa aaaaaaaaaa aaaaaaaaaa 1608

<210> 2

<211> 322

<212> PRT

<213> Haematococcus pluvialis

<400> 2

Thr Phe His Lys Pro Val Ser Gly Ala Ser Ala Leu Pro His Ile Gly
 1 5 10 15
 Pro Pro Pro His Leu His Arg Ser Phe Ala Ala Thr Thr Met Leu Ser
 20 25 30
 Lys Leu Gln Ser Ile Ser Val Lys Ala Arg Arg Val Glu Leu Ala Arg
 35 40 45
 Asp Ile Thr Arg Pro Lys Val Cys Leu His Ala Gln Arg Cys Ser Leu
 50 55 60
 Val Arg Leu Arg Val Ala Ala Pro Gln Thr Glu Glu Ala Leu Gly Thr
 65 70 75 80
 Val Gln Ala Ala Gly Ala Gly Asp Glu His Ser Ala Asp Val Ala Leu

	85		90		95
Gln Gln Leu Asp Arg Ala Ile Ala Glu Arg Arg Ala Arg Arg Lys Arg	100		105		110
Glu Gln Leu Ser Tyr Gln Ala Ala Ala Ile Ala Ala Ser Ile Gly Val	115		120		125
Ser Gly Ile Ala Ile Phe Ala Thr Tyr Leu Arg Phe Ala Met His Met	130		135		140
Thr Val Gly Gly Ala Val Pro Trp Gly Glu Val Ala Gly Thr Leu Leu	145		150		155
					160
Leu Val Val Gly Gly Ala Leu Gly Met Glu Met Tyr Ala Arg Tyr Ala	165		170		175
His Lys Ala Ile Trp His Glu Ser Pro Leu Gly Trp Leu Leu His Lys	180		185		190
Ser His His Thr Pro Arg Thr Gly Pro Phe Glu Ala Asn Asp Leu Phe	195		200		205
Ala Ile Ile Asn Gly Leu Pro Ala Met Leu Leu Cys Thr Phe Gly Phe	210		215		220
Trp Leu Pro Asn Val Leu Gly Ala Ala Cys Phe Gly Ala Gly Leu Gly	225		230		235
					240
Ile Thr Leu Tyr Gly Met Ala Tyr Met Phe Val His Asp Gly Leu Val	245		250		255
His Arg Arg Phe Pro Thr Gly Pro Ile Ala Gly Leu Pro Tyr Met Lys	260		265		270
Arg Leu Thr Val Ala His Gln Leu His His Ser Gly Lys Tyr Gly Gly	275		280		285
Ala Pro Trp Gly Met Phe Leu Gly Pro Gln Glu Leu Gln His Ile Pro	290		295		300
Gly Ala Ala Glu Glu Val Glu Arg Leu Val Leu Glu Leu Asp Trp Ser	305		310		315
					320
Lys Arg					

We claim:

1. A protein which has an enzymatic activity for converting
5 β -carotene into zeaxanthin or canthaxanthin into astaxanthin,
comprising the amino acid sequence SEQ ID NO. 2 or a sequence
derived from this sequence by substitution, insertion or
deletion of amino acids and having a homology of at least 50%
at the amino acid level with the sequence SEQ ID NO.2.
- 10 2. A protein as claimed in claim 1, wherein the protein has an
enzymatic activity for converting β -carotene into zeaxanthin
and an enzymatic activity for converting canthaxanthin into
astaxanthin.
- 15 3. A nucleic acid encoding a protein as claimed in either of
claims 1 or 2.
4. A nucleic acid as claimed in claim 3, which consists of the
20 sequence depicted in SEQ ID NO. 1.
5. A nucleic acid construct comprising a nucleic acid as claimed
in claim 3 or 4 which is functionally linked to one or more
regulation signals to increase gene expression.
- 25 6. A nucleic acid construct as claimed in claim 5, wherein the
nucleic acid as claimed in claim 3 or 4 is inserted into a
vector which is suitable for expression of the gene in a
prokaryotic or eukaryotic organism.
- 30 7. A genetically manipulated organism, where the gene expression
of a nucleic acid as claimed in claim 3 or 4

in the case where the initial organism comprises a nucleic
35 acid as claimed in claim 3 or 4 is increased by comparison
with a wild type or

in the case where the initial organism does not comprise a
nucleic acid as claimed in claim 3 or 4 is caused by the
40 genetic manipulation.
8. A genetically manipulated organism as claimed in claim 7,
whose carotenoid metabolism is different from that of the
wild type.

9. A genetically manipulated organism, as claimed in claim 7 or 8, wherein the organism used is a eukaryotic organism.
10. A genetically manipulated organism as claimed in claim 9,
5 wherein a plant is used as eukaryotic organism.
11. A process for producing genetically manipulated organisms as claimed in any of claims 7 to 10, which comprises introducing a nucleic acid as claimed in claim 3 or 4 or a nucleic acid
10 construct as claimed in claim 5 or 6 into the genome of the initial organism.
12. The use of the nucleic acid as claimed in claim 3 or 4 for producing genetically manipulated organisms.
15
13. A process for the preparation of xanthophyll derivatives, which comprises converting a β -ionone structural element into a 3-hydroxy- β -ionone structural element and/or a 4-keto- β -ionone structural element into a 3-hydroxy-4-keto- β -ionone
20 structural element in the presence of a protein as claimed in either of claims 1 or 2.
14. A process for preparing xanthophyll derivatives as claimed in claim 13, wherein an organism as claimed in any of claims 7
25 to 10 is cultivated, the organism is harvested, and the xanthophyll derivatives are then isolated from the organism.
15. The use of the protein as claimed in claim 1 or 2 for preparing xanthophyll derivatives.
30
16. The use of the nucleic acid as claimed in claim 3 or 4 for preparing xanthophyll derivatives.

35

40

45




Fig. 1




Fig. 2

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)